

Analysis of Rainfall and Temperature Trends in Bahati Sub-County, Kenya

Jackson K. Koimbori^{1,4,*}, Chris A. Shisanya², Shadrack K. Murimi³ & Rickard Petterson⁴

¹Department of Geography, Kenyatta University
Nairobi, Kenya

²Department of Geography, Kenyatta University
Nairobi, Kenya

³Department of Geography, Kenyatta University
Nairobi, Kenya

⁴Department of Earth Sciences, Uppsala Universitet
Uppsala, Sweden

*Corresponding author's email: Jackson_kinyanjui [AT] yahoo.co.uk

ABSTRACT----- *This study analyzed the climate variability trends in Bahati Sub-County, Kenya for the period 1985 to 2015. The climate parameters used included rainfall onset, rainfall cessations, seasonal rain, annual rainfall, annual mean temperature, maximum and minimum temperature. Data on rainfall and temperature was collected from Nakuru Meteorological Station. The strength and direction of the relationship between rainfall and temperature was determined using SPSS software version 23 (correlation analysis) and Microsoft Office Excel 2010 to generate frequency tables, pie charts, graphs and moving averages. The findings revealed that there is a significant, negative and strong relationship between annual rainfall trend and maximum temperature ($r = -0.716$, $p = 0.000$). However, the study findings also revealed that the relationship between annual rainfall and minimum temperature had an insignificant, weak and negative relationship ($r = 0.151$, $p = 0.418$).*

Keywords----- climate variability, rainfall, temperature

1. INTRODUCTION

The global surface temperatures in the last three decades and past century have warmed up by 0.6°C and 0.8°C respectively [4]. In 1992 the International Political Response to Climate Change (IPRCC) began with the ratification of the United Nation Framework Convention on Climate Change (UNFCCC). Stabilizing the greenhouse gases concentration in the atmosphere at a threshold that would avert severe anthropogenic interference with the present climatic systems in our world was the main purpose of the convention. This led to the signing of the Kyoto protocol of 1997 committing countries towards taking up climate action. According to the National Climate Change Action Plan (NCCAP), 2013 report, climate change can be monitored through measurement of the climatic parameters like wind, pressure, temperature and rainfall among others [3]. Climate change research studies have predicted an increase of precipitation and temperature throughout the year with an annual average of 4.2mm and +2.8°C per month respectively over tropical Africa. In Kenya data from National Climate Change Response Strategy (NCCRS) in 2010 indicated that the country's day and night temperatures have increased by as much as 2.1°C and 2.9°C respectively in western part of Kenya in the last twenty years. Meanwhile in central Kenya, which includes Nairobi the night and day temperatures have risen by 2.0°C and 0.7°C respectively. In the southern-eastern region, which includes Kenya's food basket, the rift valley temperatures have risen by 1°C. In Nairobi both maximum and minimum temperatures have shown an increasing trend for the period 1966 to 1999 [9]. Climate change and variability in Kenya has led the Government of Kenya to commit approximately Kenya Shillings (KES) 37 billion, while development partners have committed KES 194 billion between 2005 and 2015 to programs they have earmarked as having a major impact on climate change components [3]. The residents of Bahati Sub-County, Kenya have been experiencing increasing temperatures and variations in rainfall amounts. This research study therefore sought to analyse the rainfall and temperature trends in Bahati Sub-County, Kenya for the period 1985 to 2015 with the aim of providing the locals with a better understanding of the rainfall and temperature trends.

2. METHODOLOGY

2.1 Study Area

The study was carried out in Bahati Sub-County which is located within Nakuru County, Kenya (Fig. 1). It is made up of 5 wards of which 4 wards will be studied; Dundori ward, Kabatini ward, Kiamaina ward and Bahati ward, since Lanet/Umoja is not an agricultural area. The greater Nakuru County is in Rift valley province and occupies an area of 7,242.3km². It is located between longitudes 35°28' and 35° east and Latitude 0°13' and 10°10' South at an altitude of about 1912 meters above sea level. Bahati Sub-County has a population of 141,352 covering an area of 375.40km² [7]. Nakuru County has predictable weather patterns with temperatures ranging between 10°C during the cold months (July and August) and 20°C during the hot months (January to March). Bahati Sub-County receives between 700mm and 1200mm of rainfall annually, with average annual rainfall being an approximated 960mm. Nakuru County which covers Bahati Sub-County has two rainy seasons; March, April, May (long rains) and October, November, December (short rains) [3]. The soils are complex due to influence by variations in relief, climate and underlying rock types. The major soils fall into three categories; Latosolic soils, Planosolic soils and Alluvial soils [3].




Figure 1. A Map of Bahati sub-county showing study site.

2.2 Climate Data

A span of the most recent thirty year records from 1985 to 2015 was collected from Nakuru Meteorological Station. The climate data was a representative of the most recent average climate in the study region and the 30 years' span was sufficient duration to encompass a range of significant weather anomalies depicting monthly rainfall and temperature trends in Bahati Sub-County, Nakuru County. The rainfall and temperature characteristics for Bahati Sub-County were recorded in a summary check sheet and this included rainfall onset, rainfall cessations, seasonal rain, annual rainfall, seasonal maximum and minimum temperature.

2.3 Data Analysis

The collected data was analysed using descriptive and inferential statistics. Descriptive statistics included measures of central tendency and measures of dispersion. The mean rainfall and temperature for the period 1985 to 2015 was determined from Nakuru Meteorological Station. Consequently, statistics on the inter-annual standard deviation and variance of rainfall and temperature for the same period were calculated.

3. RESULTS

3.1 Rainfall Trends

3.1.1 Annual Rainfall Trend

The results in Fig. 2, shows that the annual rainfall amount has reduced between 1985 and 2015 in Bahati Sub-County. High amounts of rainfall are noticed in the years 1988 (1244mm), 1992 (980mm), 1997 (1094mm), 2001 (1130mm), 2002 (1084mm), 2003 (1138mm), 2007 (1217mm), 2010 (1436mm) and 2013 (1185mm), while low amounts of rainfall are noticed in the years 1987 (697mm), 1993 (704mm), 2000 (610mm) and 2009 (705mm). The rainfall variability shown in the results above supports the argument that the ASALs in Kenya have been impacted negatively by high rainfall variability [10, 11].




Figure 2: Annual Rainfall Trend for Bahati Sub-County

3.1.2 Seasonal Rainfall Trends

The extreme events were observed both during the long and short rainfall season. During the long rainfall season high amounts of rainfall were recorded during the years 1985 (505mm), 1988 (575mm), 1990 (487mm), 2003 (464mm), 2010 (585mm), and 2012 (453mm), while low amounts of rainfall were recorded in 1993 (125mm), 1996 (199mm), 2000 (69mm), and 2008 (179mm) (Fig. 3). Further, during the short rainfall season high amounts of rainfall were recorded during the years 1989 (347mm), 1997 (464mm), 2002 (384mm), 2006 (352mm), 2008 (311mm) and 2015 (315mm), while low amounts of rainfall were recorded in 1985 (106mm), 1993 (126mm) and 2005 (112mm) (Fig. 3). According to a 2007 report by Intergovernmental Panel on Climate Change (IPCC), changes in rainfall pattern in East Africa is as a result of the El Niño phenomena [6] and this supports our findings on the seasonal rainfall pattern fluctuations in Bahati Sub-County.




Figure 3: Seasonal Rainfall Trend for Bahati Sub-County

3.2 Rainfall Variability

The annual rainfall variability characteristics for Bahati Sub-County (1985 to 2015) were computed when annual rainfall anomalies were presented in the graph as shown in Fig. 4. Annual rainfall variability ranges from -350.40mm in 2000 to +475.80mm in 2010 as shown in Fig. 4. The highest rainfall anomalies were recorded in years 1987 (-263.2mm), 1988 (+283.2mm), 1993 (-256.5mm), 1999 (-303.6mm), 2000 (-350.4mm), 2007 (+256.9mm), 2009 (-255.1mm), 2010 (+475.8mm) and 2013 (+224.4mm).




Figure 4: Annual Rainfall Variations for Bahati Sub-County

3.2.1 Seasonal Rainfall Variations

The variability in seasonal rainfall (long and short), and occurrence of extreme events have effects on maize yield in the area exposing small scale farmers to climate change vulnerability. This collaborates with various studies that have shown that changes in seasonal rainfall patterns have a negative effect on rain fed agriculture [2,5]. Rainfall variability is significant in the long rain season trend and ranges from (-253.17mm) in 2000 to (+263.53mm) in 2010 as shown in Fig. 5. Rainfall variability is also significant in the short rain season trend and ranges from (- 111.08mm) in 1985 to (+247.31mm) in 1997 as shown in Fig. 5. During the long rains (MAM) the highest rainfall variations were recorded in 1985 (+182.8mm), 1988 (+253.0mm), 1993 (-196.6mm), 2000 (-253.2mm) and 2010 (+263.5mm), while during the short rains (OND) the highest rainfall variations were recorded in 1985(-111.08mm), 1989 (+130.7mm), 1997 (+247.3mm), 2002 (+167.0mm), 2005 (-104.6mm) and 2006 (+134.8mm).




Figure 5: Seasonal Rainfall Variations for Bahati Sub-County

3.3 Seasonal Rainfall Onset and Cessation

The results in Table 1 shows that onset month for Long rain season has varied, alternating between March and April. 18 times representing 58.1% the onset was in March and 13 times representing 41.9% the onset was in April. Cessation month has also varied alternating between June, July and August. When farmers are informed on the onset date they plan on when to prepare their land and acquire the necessary inputs. This finding supports the earlier findings by [1] that the onset time for long term rainfall is normally used by farmers to determine the time they clear and prepare the land for planting. The peak months' range from March, April and May. The results in Table 1 shows that the onset month for the short rain season has varied, alternating between October, November and December. 23 times representing 74.2% the onset was in October, 7 times representing 22.6% the onset was in November and 1 time representing 3.2% the onset was in December. Cessation month has also varied alternating between November, December and January. The peak months' range from October, November and December. The results found that seasonal rainfall change is a challenge to Bahati Sub-County farmers and they need to be updated at all times to enable them plant on time.

Table 1: Summary of Seasonal Rainfall Characteristics for Bahati Sub-County

Year	Peak Month of Rainfall During the Long Rain Season	Onset Month of Rainfall During the Long Rain Season	Cessation Month of Rainfall During the Long Rain Season	Peak Month of Rainfall During the Short Rain Season	Onset Month of Rainfall During the Short Rain Season	Cessation Month of Rainfall During the Short Rain Season
1985	April	April	July	November	November	December
1986	April	April	July	December	December	January
1987	May	April	July	November	October	December
1988	May	March	June	October	October	January
1989	April	April	June	October	October	January
1990	April	March	June	October	October	December
1991	March	March	June	November	November	December
1992	April	April	June	October	October	November
1993	May	April	July	November	October	December
1994	May	March	July	November	November	December
1995	March	March	July	October	October	December
1996	May	March	September	November	November	December
1997	April	April	June	November	October	December
1998	May	April	July	November	November	December
1999	March	March	June	December	November	January
2000	April	April	September	November	October	December
2001	April	March	September	October	October	December
2002	April	March	June	December	October	January
2003	May	March	June	October	October	December
2004	April	March	June	December	October	January
2005	May	March	July	October	October	December
2006	May	March	June	November	October	January
2007	May	April	June	October	October	December
2008	March	March	June	October	October	December
2009	May	April	June	December	October	January
2010	March	March	June	October	October	December
2011	May	March	June	November	October	December
2012	April	April	June	October	October	January
2013	April	March	August	December	November	January
2014	May	March	June	October	October	January
2015	April	April	June	November	October	January

***.* Correlation is significant at the 0.01 level (2-tailed).

3.4 Temperature Trends

3.4.1 Annual Average Temperature Trend

Temperature being the other key indicator of climate variability, this study also sought to determine the temperature variations of the area for the last 30 years (1985 to 2015). The mean annual temperature in the area has generally been increasing since 1985 to 2015. The lowest average temperature recorded in Bahati Sub-County was 17.7°C in 1989, while the highest temperature recorded was 19.7°C in 2009, (Fig. 6) and this coincides with the years 1987, 2000 and 2009 when Kenya experienced the worst droughts. According to [9], seasonal temperature changes observed could alter the growing, planting and harvesting time for agricultural production.




Figure 6: Annual Average Temperature Trend for Bahati Sub-County

3.4.2 Annual Average Maximum Temperature Trend

The maximum average temperature for the Bahati Sub-County (1985 to 2015) was 25.8°C, while the highest maximum annual temperature was recorded in 1987 and 2009 at 27.0°C and the lowest in 1989 at 24.8°C as shown in Fig. 7. The Trend line equation shows that Bahati-Sub-County has been experiencing a slightly gradual increase of 0.008°C annually in the maximum temperature for the period 1985 to 2015.




Figure 7: Annual Average Maximum Temperature for Bahati Sub-County

3.4.3 Annual Average Minimum Temperature Trend

The minimum average temperature for Bahati Sub-County for the period 1985 to 2015 was recorded at 11.6°C. The lowest temperature recorded was in 1986 at 9.6°C and the highest minimum temperature recorded was in 2010 at 12.7°C as shown in Fig. 8. The trend line equation shows that there has been a gradual increase of 0.075°C in minimum temperature for Bahati Sub-County for the period 1985 to 2015.




Figure 8: Annual Average Minimum Temperature for Bahati Sub-County

3.5 Temperature Variability

3.5.1 Annual Average Temperature Variations

Annual average temperature variability characteristics for Bahati Sub-County showing the anomalies for the period 1985 to 2015 was presented in a graph as shown in Fig. 9. Annual average temperature variability ranged from 17.7°C in 1989 to 19.7°C in 2009 (Fig. 6). The highest average temperature anomalies were recorded in the years 1986(-0.88°C), 1989(-1.0°C), 2009(+0.98°C), and 2015(+0.91°C) as shown in Fig. 9.




Figure 9: Annual Average Temperature Variations for Bahati Sub-County

3.5.2 Annual Average (Maximum and Minimum) Temperature variations from the Mean.

Maximum and minimum temperature variability for Bahati Sub-County for the period 1985 to 2015 was recorded as significant with maximum temperature variations peaks noted during the years 1987(+1.2°C), 1988(-0.7°C), 1989(-1.1°C), and 2009(+1.2°C) as shown in Fig. 10. During the minimum temperature variations, high temperature variations were recorded during the years 1985(-1.7°C), 1986(-2.1°C) and in 2010(+1.1°C).




Figure 10: Annual Average (Max. and Min.) Temperature Variations for Bahati Sub-County

Table 2: Summary of Recorded Rainfall and Temperature Data for Bahati Sub-County (1985 to 2015)

Annual	Rainfall (mm)	Temperature (°C)
Maximum	1436.2mm	27.0°C
Minimum	580.9mm	9.6°C
Mean	960.4mm	18.7°C

Source: Kenya Meteorological Department (1985 to 2015)

3.6 Correlation Analysis

3.6.1 Correlations of Annual Rainfall, Maximum Annual Temperature and Minimum Temperature

The annual rainfall trend, maximum temperature trend and minimum temperature trend as shown on Table 4 were correlated to determine the significance of their relationship. When Pearson's r is close to 1, this means that there is a strong relationship between the two variables and that changes in one variable are strongly correlated with changes in the second variable. When Pearson's r is close to 0 this means that there is a weak relationship between the two variables and that the changes in one variable are not correlated with changes in the second variable. From the findings on Table 4, Pearson's ($r = -0.716$, $p = 0.000$) is close to -1 showing that there is a significant strong and negative linear relationship between annual rainfall amount and maximum temperature. Pearson's r ($r = 0.151$, $p = 0.418$) recorded between annual rainfall and minimum temperature indicated an insignificant, weak and negative relationship between the two variables. A positive Pearson's r indicates that as one variable increases, the second variable also increase in value or vice versa.

Table 3: Annual Rainfall, Maximum Annual Temperature and Minimum Temperature (1985 to 2015)

Years	Annual Rainfall Amount (mm)	Maximum Annual Temperature Amount (°C)	Minimum Annual Temperature Amount (°C)
1985	987	25.9	9.9
1986	934.2	26.1	9.6
1987	697.2	27.0	10.9
1988	1243.6	25.1	11.1
1989	1145.4	24.8	10.6
1990	978.6	25.2	10.6
1991	768.4	26.0	10.6
1992	979.8	25.5	10.9
1993	703.9	25.8	10.9
1994	790.4	25.6	11.4
1995	917.8	25.8	11.3
1996	933.1	25.6	11.2
1997	1094.2	25.6	11.5
1998	988.4	25.3	12.5
1999	656.8	26.0	11.6
2000	610	26.6	12.0
2001	1129.8	25.4	12.0
2002	1083.7	25.8	11.9
2003	1137.6	25.6	12.0
2004	891.7	25.8	11.8

2005	884.5	26.1	11.5
2006	971.5	25.7	12.4
2007	1217.3	25.2	12.0
2008	825.8	25.7	12.1
2009	705.33	27.0	12.4
2010	1436.2	25.3	12.7
2011	1109.9	25.8	12.2
2012	1134	25.6	11.9
2013	1184.8	25.6	12.2
2014	814.8	26.1	12.4
2015	816.7	26.7	12.5

Table 4: Summary of Correlations between Annual Rainfall, Maximum Annual Temperature and Minimum Temperature for Bahati Sub-County

		Rainfall (mm)	Maximum Temperature (°C)	Minimum Temperature (°C)
Rainfall (mm)	Pearson Correlation	1	-.716**	.151
	Sig. (2-tailed)		.000	.418
	N	31	31	31
Maximum Temperature (°C)	Pearson Correlation	-.716**	1	.081
	Sig. (2-tailed)	.000		.663
	N	31	31	31
Minimum Temperature (°C)	Pearson Correlation	.151	.081	1
	Sig. (2-tailed)	.418	.663	
	N	31	31	31

** . Correlation is significant at the 0.01 level (2-tailed).

4. DISCUSSION

4.1 Climate Variability

The study found out that the annual rainfall had reduced between 1985 and 2015 in Bahati Sub-County. High amounts of rainfall were noticed in the years 1988 (1244mm), 1992 (980mm), 1997 (1094mm), 2001 (1130mm), 2002 (1084mm), 2003 (1138mm), 2007 (1217mm), 2010 (1436mm) and 2013 (1185mm), while low amounts of rainfall are noticed in the years 1987 (697mm), 1993 (704mm), 2000 (610mm) and 2009 (705mm). Seasonal rainfall variability was

found to be significant in the long and short rain season trend and ranged from -253.17mm (2000) to +263.53mm (2010) and from -111.08mm (1985) to +247.31mm (1997) respectively. The mean annual temperature in the area has generally been increasing since 1985 to 2015. The lowest average temperature recorded in Bahati Sub-County was 17.7°C in 1989, while the highest temperature recorded was 19.7°C in 2009, (Fig. 6). The trend line equation showed that Bahati-Sub-County has been experiencing a slightly gradual increase of 0.008°C annually in the maximum temperature for the period 1985 to 2015. While the highest maximum annual temperature was recorded in 1987 and 2009 at 27.0°C and the lowest in 1989 at 24.8°C as shown in Fig. 7. The trend line equation showed that there has been a gradual increase of 0.075°C in minimum temperature for Bahati Sub-County for the period 1985 to 2015. The lowest temperature recorded was in 1986 at 9.6°C and the highest minimum temperature recorded was in 2010 at 12.7°C as shown in Fig. 8.

5. CONCLUSIONS

The main objective of the study was to analyze the climate variability trends in Bahati Sub-County, Kenya for the period 1985 to 2015. The study found out that the annual rainfall amounts had decreased over the years, while maximum and minimum temperatures over the years have gradually been increasing. The correlation analysis revealed that there is a significant, negative and strong relationship between annual rainfall trend and maximum temperature ($r = -0.716$, $p = 0.000$). However, the study findings also revealed that the relationship between annual rainfall and minimum temperature had an insignificant, weak and negative relationship ($r = 0.151$, $p = 0.418$). Furthermore, the study findings revealed that there has been a shift in the onset and cessation of rainfall. The onset month for long rain season has been alternating from March to April while the cessation month has been alternating from June to August. The onset month for the short rainy season has varied alternating between October to December, while the Cessation month has also varied alternating between November to January.

6. ACKNOWLEDGEMENTS

I would like to acknowledge the assistance and support accorded to me by the Kenya Meteorological Department, Department of Geography in Kenyatta University and Department of Earth Sciences in Uppsala Universitet.

8. REFERENCES

- [1] Adeniyi M., Ogunisola O., Nymphas E, and Oladiran O. (2009). Food Security Measures During Uncertain Climatic Conditions in Nigeria. University of Ibadan, Ibadan, Nigeria. *African Journal of Food Agriculture Nutrition and Development*, 9:652-677.
- [2] Bals, C., Harmeling, S. and Windfuhr, M., (2008). *Climate change, food security and the right to adequate food*. Stuttgart, Germany: Diakonie Katastrophenhilfe, Brot fuer die Welt and German Watch.
- [3] Government of Kenya, (2013). *National Climate Change Action Plan 2013 -2017*. Government printers, Nairobi, Kenya.
- [4] Hansen, J., Sato, M., Ruedy, R., Lo, K., Lea, D.W. and Medina-Elizade, M. (2006). *Global temperature change*. Proceedings of the National Academy of Sciences of the United States of America, 103(39).
- [5] IITA. (2004). *Annual report for Ibadan, Nigeria*. International Institute for Tropical Agriculture.
- [6] IPCC. (2007). *Climate Change 2007: Impacts, adaptation and vulnerability: Contribution of Working Group II to the fourth assessment report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK.
- [7] KNBS. (2009). *Kenya Population and Housing Census. Population Distribution by Age, Sex and Administrative Units*, IC. Kenya National Bureau of Statistics, Nairobi: Kenya.
- [8] Makokha, G. L. and Shisanya, C.A. (2010). Trends in Mean Annual Minimum and Maximum Near Surface Temperature in Nairobi City, Kenya. *Advances in Meteorology*, 2010: 6.
- [9] Mark, W., Mandy, E., Gary, Y., Lan, B., Saleemul, H., and Rowena, V. (2008). *Climate Change and Agriculture. Threats and Opportunities*. Federal Ministry for Economic Cooperation and Development, Germany.
- [10] Omoyo, N.N., Wakhungu, J. and Oteng'i, S. (2015). *Effects of climate variability on maize yield in the arid and semi-arid lands of lower eastern Kenya*. *Agriculture and Food Security* 4:8.
- [11] Shisanya, C.A., Recha, C. and Anyamba, A. (2011). Rainfall Variability and Its Impact on Normalized Difference Vegetation Index in Arid and Semi-Arid Lands of Kenya. *International Journal of Geosciences*, 2, 36-47